

breast cancer and is based on how well it is working and what side effects (5). Chemo drugs can cause side effects such as hair loss and nail changes, mouth sores, loss of appetite or increased appetite and nausea and vomiting, increased susceptibility to infection, easy bruising, depending on the type and dose of drugs given, and the length of treatment (4).

Oncology Health Care Providers (HCPs) generally assume that patients with cancer will adhere to treatment recommendations because of the seriousness of a cancer diagnosis, however, reports in the literature have demonstrated adherence levels as low as 20% (6, 7). Reasons for non-adherence are complex in most situations (8). Factors that have been frequently associated with non-adherence to recommended medical therapies include individual patient characteristics, features of the disease and the treatment regimen, and aspects of the medical care system (9). Failure to adherence of medication is a serious problem which does not only affect the patient but also the healthcare system (10). Medication non-adherence leads to substantial worsening of the disease, death and increased health cost and as such health care professionals such as nurses, doctors and pharmacist have a significant role in their daily practice to improve patient adherence to medication (2).

Nigerian health care system faces public health challenges in the form of breast cancer. The issue of non-adherence to chemotherapeutic agents among women with breast cancer has been cited by the World Health Organization (2008) as the single most important modifiable factor that can compromise treatment outcomes. Non-adherence is dangerous and can lead the practitioner to change the dose or prescribe a different agent because of apparent non-responsiveness or unexpected adverse effects. It can also result in unnecessary diagnostic testing, changes in dose or therapeutic regimen and hospitalizations (11). It has been observed at the University of Nigeria Teaching Hospital, Enugu that there is an increase in oncologist visit by cancer survivors due to a relapse in the

disease condition. Increased hospitalization rates, longer hospital stays and decreased patient satisfaction are all possible outcomes of non-adherence to chemotherapy. This study is thus aimed at identifying factors that may impede the success of chemotherapy as these will enable Oncology health care providers to plan accurately for the treatment regimen and to improve the quality of health services provided. The specific objectives of this study are to ascertain the patient related factors, therapy related factors and health care system factors that influence non-adherence to chemotherapy among breast cancer survivors at UNTH, Enugu.

Methods

This is a cross sectional descriptive survey design study that carried out in University of Nigeria Teaching Hospital (UNTH) Enugu –Nigeria. The population for the study comprised of 100 breast cancer survivors currently receiving chemotherapy at oncology clinic of UNTH, Enugu - Nigeria. The sampling method used was census due to the small population. The entire population was used utilizing inclusion criteria of willingness of respondents to participate, and respondents on chemotherapy.

The instrument used in obtaining information for this study was the Breast Cancer questionnaire which was constructed in line with the research objectives. The Breast cancer questionnaire consisted of four sections. Section A which address the respondents' socio-demographic data and section B that determine the patient related factors that influence the non-adherence to chemotherapy among breast cancer patients at UNTH, Section C that determine the therapy related factors that influence non-adherence to chemotherapy among breast cancer patients at UNTH and section D which assess the health care system factors that influence non-adherence to chemotherapy among breast cancer patients at UNTH.

Table 1. Socio-demographic profile of the respondents (n=100)

Variables		N	%
Age distribution in years	20-29	11	11.0
	30-39	32	32.0
	40-49	32	32.0
	50-59	18	18.0
	60 and above	7	7.0
Highest level of education	No formal education	4	4.0
	Primary education	7	7.0
	Secondary education	29	29.0
	Tertiary education	60	60.0
Religion	Christianity	94	94.0
	Islam	6	6.0
Occupation	Housewife	10	10.0
	Farmer	4	4.0
	Trader/ business	16	16.0
	Civil servant	41	41.0
	Self-employed	21	21.0
	Not employed	8	8.0
Marital status	Married	70	70.0
	Single	26	26.0
	Widowed	4	4.0

Table 2. Chemotherapy regimen characteristics among the respondents (n=100)

Variable questions	N	%	
When diagnosed of breast cancer	Less than four months	22	22.0
	More than six months	36	36.0
	Over a year	21	21.0
	Over two years	21	21.0
When planned for chemotherapy	<Less than four days after diagnosis	7	7.0
	A week after diagnosis	21	21.0
	Two weeks after diagnosis	12	12.0
	A month after diagnosis	25	25.0
	More than one month after diagnosis	35	35.0
Commencement of Chemotherapy	1-2 weeks after planning	20	20.0
	3-4 weeks after planning	15	15.0
	>1 month after planning but < 2months	43	43.0
	>2 months	22	22.0
What delayed the commencement of chemotherapy	Busy work schedule	8	8.0
	Financial constraint	48	48.0
	Lack of family/community support	3	3.0
	Did not believe result	6	6.0
	No response	27	27.0
Courses of chemotherapy recommended	Four	50	50.0
	Eight	21	21.0
	Twelve	10	10.0
	Sixteen	3	3.0
	More than seventeen	16	16.0
Courses of chemotherapy missed	None	21	21.0
	One	19	19.0
	twice	31	31.0
	More than two times	9	9.0

non-adherence among women taking Tamoxifen (13). These side effects according to Schneider, Hess and Gosselin, (2013) includes alopecia, leucopenia, and pain, loss of appetite or increased appetite, increased susceptibility to infection, easy bruising, thrombocytopenia, infertility, anaemia, and dermatitis. This result agrees with Adewale, Olukayode and Adesunkanmi (11), study which reported that 11% of their study sample size was unable to further bear the drug side effects. Duration of treatment was also seen to be another therapy related factor that hinders adherence to chemotherapy; this result is in contrast with Adewale, Olukayode and Adesunkanmi study which reported that 15% of their study sample size identified fear of subsequent operation. This could be associated with the worry associated with informing patient that they would receive treatment for a long time. Another factor influencing chemotherapy non-adherence is the respondents' being told to abstain from eating some foods like meat and preference to taking drugs which might be due to the non-painful effect of taking such drug. This agrees with Kelloway, Wyatt and Adlis, (2014) (14) who identified that medications with a convenient way of administration (eg, oral medication) are likely to make patients compliant compared with injections.

The result of the study revealed that prescribed drug unavailable in the hospital pharmacy 47(47.0%) and the equal identification of patient-prescriber relationship and unfavorable hour of clinic visit 40(40%) respectively influences the respondents' adherence to chemotherapies. In the present economy, the poor budgetary allocation for 2017, meagre 4.15 per cent, though a marginal improvement on the 3.73 per cent figure of 2016 as against the WHO standard for member countries to give at least 13 per cent of its minimum allocation to the health sector if they must adequately meet the healthcare needs of their citizens. The implication of this is an under stocked pharmacy in the hospital. Also, in the area of the present study, the hospital is short staffed, this possibly resulted in designation

of clinic days which may be unfavorable for patients thus the few personnel available attend to a great number of the patients on specified clinic days. This inadvertently may result in burnout on the part of the health care provider, considering the large number of patients they need to attend to, thereby creating a strain in patient-provider relationship. This is in contrast with the study of Ponnusankar (15) who discovered availability and accessibility to health care and lack of accessibility to healthcare as healthcare related factors hindering adherence to chemotherapy while Lawson et al (16) identified long waiting time for clinic visits and difficulty in getting prescriptions filled (17), unhappy or unsatisfied clinic visits (16) which all contributed to poor compliance.

The result of the study demonstrated the various factors that influence patients' non-adherence to chemotherapy. Based on the result, financial constraint and worry about outcome of treatment, medication side effects such as hair lose, loss of weight, duration of treatment and prescribed drugs not available in the hospital pharmacy and unfavorable hour of clinic visit. This implies the nurses to design various programs such as freewill donations from church organizations and the affluent members in the society as well as the government agencies to inspire the cancer survivors in adhering to therapy. Also, health education initiation on benefit of adhering to chemotherapy would be needed on the part of the nurses to foster the women intake of chemotherapy. Nurses caring for women who receive endocrine therapy for breast cancer should identify those who may be at greater risk for being non-adherent. Therefore, nurses should assess cancer survivors to determine whether they are experiencing depression or anxiety during pre-therapy and throughout the course of therapy. Nurses also should assess women to determine whether they are experiencing symptoms related to their breast cancer or its treatment. Effective management of problems with symptoms experienced at pre-therapy and throughout therapy may help women with breast cancer

maintain better adherence to endocrine therapy over time. Also, nurses need to assess whether women are experiencing financial hardships related to the cancer and its treatment and provide appropriate information and referrals so that women receive the resources they need to complete the full, prescribed course of chemotherapy. It was therefore recommended that the nurses should advocate for the patients for the federal government to re-implement the health care policy that allowed treatment free-of-charge at Nigerian Government hospitals to those with malignancies and other chronic ailments to mitigate the burden of associated financial problem and encourage patients to seek orthodox medical care. The fee for chemotherapy should also be subsidized as this would improve patient's adherence to chemotherapy. There should be cancer specific educational package with focus on knowledge about disease and treatment to be organized routinely by the health care personnel in the various health care facilities in Nigeria. More health care personnel should be made available to work in the health facility to reduce the problem of unfavorable hour of clinic visits and burnout associated with short staffing. An increase in the federal government allocation towards the health sector to reduce the problem of drug unavailability in the hospital pharmacy. The uncooperative attitude of the respondents, the small sample size which may introduce bias and limited materials on the topic.

Acknowledgement

Authors will want to appreciate the statistician for his support in the data analysis.

Conflicts of Interest

The authors declare that there are no conflicts of interest regarding the publication of this paper.

References

1. World Health Organization (WHO). Breast cancer; prevention and control. 2016

[updated 2016; cited]; Available from: <https://www.who.int/cancer/detection/breastcancer/en/>.

2. Okoronkwo I, Ejike-Okoye P, Chinweuba A, Nwaneri A. Financial barriers to utilization of screening and treatment services for breast cancer: an equity analysis in Nigeria. *Nigerian journal of clinical practice*. 2015;18(2):287-91.

3. Shulman LN, Willett W, Sievers A, Knaul FM. Breast cancer in developing countries: opportunities for improved survival. *Journal of oncology*. 2010;2010.

4. Siegel RL, Miller KD, Jemal A. Cancer statistics, 2017. *CA: a cancer journal for clinicians*. 2017;67(1):7-30.

5. Hadi MA, Hassali MA, Shafie AA, Awaisu A. Knowledge and perception of breast cancer among women of various ethnic groups in the state of Penang: a cross-sectional survey. *Medical Principles and Practice*. 2010;19(1):61-7.

6. Siqueira JLD, Morete MC. Psychological assessment of chronic pain patients: when, how and why refer? *Revista Dor*. 2014;15(1):51-4.

7. Xu S, Yang Y, Tao W, Song Y, Chen Y, Ren Y, et al. Tamoxifen adherence and its relationship to mortality in 116 men with breast cancer. *Breast Cancer Res Tr*. 2012;136(2):495-502.

8. Yap AF, Thirumoorthy T, Kwan YH. Medication adherence in the elderly. *Journal of Clinical Gerontology and Geriatrics*. 2016;7(2):64-7.

9. Sabaté E. Adherence to long-term therapies: evidence for action. Geneva, Switzerland: World Health Organization; 2003. Google Scholar. 2017.

10. Jimmy B, Jose J. Patient medication adherence: measures in daily practice. *Oman medical journal*. 2011;26(3):155.

11. Adisa AO, Gukas ID, Lawal OO, Adesunkanmi ARK. Breast Cancer in Nigeria: Is Non-Adherence to Chemotherapy Schedules a Major Factor in the Reported Poor Treatment Outcome? *The breast journal*. 2010;16(2):206-7.

12. Anyanwu SN, Egwuonwu OA, Ihekwoaba EC. Acceptance and adherence to treatment among breast cancer patients in Eastern Nigeria. *The Breast*. 2011;20:S51-S3.

13. Atkins L, Fallowfield L. Intentional and non-intentional non-adherence to medication amongst breast cancer patients. *Eur J Cancer*. 2006;42(14):2271-6.

14. Kelloway JS, Wyatt RA, Adlis SA. Comparison of patients' compliance with

prescribed oral and inhaled asthma medications. *Arch Intern Med.* 1994;154(12):1349-52.

15. Ponnusankar S, Surulivelrajan M, Anandamoorthy N, Suresh B. Assessment of impact of medication counseling on patients' medication knowledge and compliance in an outpatient clinic in South India. *Patient education and counseling.* 2004;54(1):55-60.

16. Lawson VL, Lyne PA, Harvey JN, Bundy CE. Understanding why people with

type 1 diabetes do not attend for specialist advice: a qualitative analysis of the views of people with insulin-dependent diabetes who do not attend diabetes clinic. *Journal of health psychology.* 2005;10(3):409-23.

17. Vlasnik JJ, Aliotta SL, DeLor B. Medication adherence: factors influencing compliance with prescribed medication plans. *The Case Manager.* 2005;16(2):47-51.