



Original Article

Newly hired nurses' transitional challenges and coping with caring for schizophrenia patients in a psychiatric training hospital

Reynold Culimay Padagas^{1*}, John Vinnie Panuelos², Kim Franco Odoya², Rhencel Cabardo², Vincent Pierre Louise Castro², Mary Ann Manio², Alexandrae Nabablit², Mikaela Angela Papilla²

¹College of Nursing and Health Sciences, Jose Rizal University, Mandaluyong, Philippines

²Our Lady of Guadalupe Colleges, Mandaluyong, Philippines

ARTICLE INFO

Received 27 April 2020
Accepted 24 July 2020

Available online at:
<http://npt.tums.ac.ir>

Key words:

transitional challenges;
coping mechanisms;
psychiatric nurse;
qualitative research

ABSTRACT

Background & Aim: This study aimed to explore the transitional challenges of newly hired psychiatric nurses in the care of clients with schizophrenia and the coping mechanisms that they use to overcome their transitional challenges.

Methods & Materials: This descriptive-qualitative study interviewed six purposively selected newly hired psychiatric nurses caring for patients with schizophrenia who work for about six months to two years in a Psychiatric Training Hospital in the Philippines. An interview protocol was used to obtain data with strict observance of relevant ethical considerations. In determining the emerging themes and subthemes from the transcribed data with trustworthiness, thematic analysis was employed.

Results: Two major prior codes were explored, namely, transitional challenges and coping mechanisms. On the one hand, five subthemes emerged under the transitional challenges that include emotional challenges, need for professional growth, nurses' recognition of the inability to patient care management, need support from more experienced colleagues. Experience/feel the shock in the encounter of people with a mental health condition. On the other hand, four subthemes were unearthed under coping mechanisms that include emotional responses, professional coping strategies (i.e., reliance on the adequate support system, establishing the professional nurse-patient relationship, call to professional advancement, and commitment to evidence-based practice), and institutional coping strategies.

Conclusion: While newly hired nurses experience tremendous transitional challenges, their coping mechanisms continuously emerge and develop. The nursing administration needs to foster sustainable mentoring and resilience programs essential in aiding newly hired nurses' adaptation and adjustment to the professional work environment.

Introduction

Mental health nurses work in challenging and potentially high-stress settings (1), and this is more shuddering among newly hired nurses in psychiatric facilities. When psychiatric nurses are beginning to be involved in the hospital setting and realize that they are not fully equipped and competent when adjusting to and exploring a new environment, they need to be oriented and upskilled. Nursing and hospital orientation program should be improved and empowered to prepare newly hired nurses or novice nurses for the psychiatric setting (1).

The transitional process involves challenges that newly hired nurses need to endure and overcome. The transition category for this study is the situational transition (2) based on the Transitions Theory of Meleis. Situational transition is defined as a transition in various educational and professional roles, which includes the new graduate registered nurses' transition and mastery of the professional roles (2). Transforming from student nurse to registered nurse is intertwined with many challenges in terms of the environment, student nurses are used to (2). New graduate nurses, as "people in transition," need the assistance of the facilitator to minimize potential risks, guide professional development, and contribute to nursing role mastery (2). The awareness of problems

*Corresponding Author: Reynold Culimay Padagas, College of Nursing and Health Sciences, Jose Rizal University, Mandaluyong, Philippines.
E-mail: reynold.padagas@gmail.com

such as lacking a sense of security, professional competency, and having little understanding of mental illness is needed to help nurses in providing appropriate care for clients (3).

Insufficient professional skills and knowledge, ability to cope with stress and utilize emotions, and support from colleagues and team, and a lack of positive expectations about the future and emotional preparation for the job, are among the many direct or indirect difficult situations nurses encounter (4). Understanding their limitations, psychiatric nurses learn to adapt to their new work using various coping mechanisms, and these mechanisms speak about their resilience and adaptation. For instance, some psychiatric nurses used problem-focused coping to deal with patients with aggressive behaviors. The nurses managed the aggressive behavior by talking to the patient calmly, lovingly, and by leaving the patient alone (5). In a study, results show that self-respect and endurance (controllability and positive thinking) had significant positive relationships with resilience, and nurses with high level of resilience had lower levels of depression and exhaustion (emotional exhaustion and desensitization) (6). According to another study, a complex interplay between intrapersonal, interpersonal, and organizational factors influence transition (7).

In this regard, the study focused on uncovering the transitional challenges and coping mechanisms of newly hired psychiatric nurses caring for schizophrenia patients in a psychiatric facility in the Philippines.

Methods

Design

This study employed descriptive-qualitative design because it does not intend to penetrate the data in any interpretive depth and opts to present comprehensive summaries of phenomena (8). Moreover,

the design tends to be eclectic and based on the premises of the naturalistic inquiry (8).

Participants

The study involved six newly hired psychiatric nurses with six months to two years of clinical experience in a Psychiatric Training Hospital (PTH). Six participants (9) are already enough to reach data saturation. The recommended number of participants range in a number of five to twenty-five (10) or at least six for a qualitative study (11). Only the novice to advance beginner nurses (12) were identified having experience as from the first day of their job as a nurse to 2 years of experience in the field; hence, purposive sampling was used.

Data collection

An interview protocol was used because it serves to remind the researchers to relay essential bits of information to the interviewees (13). The interview protocol focused on specific questions, such as “Describe your transitional stage as a newly hired psychiatric nurse” and “What coping mechanisms do you use in coping with your challenges related to your transition?”. The researchers conducted a simulation on how to facilitate face-to-face interviews prior to actual interviews.

The interviews were audiotaped, and the recorded conversations were transcribed. The researchers referred back the transcribed data to the participants for triangulation purposes. The researchers were peer debriefed regarding data analysis by an expert with rich experience in psychiatric nursing and qualitative research. Each interview lasted for at least 30 to 45 minutes. The study was conducted in 2018.

Data analysis

In this study, thematic analysis was conducted after transcriptions were made through the process of coding in six phases to create and establish meaningful patterns. The phases included familiarization of data, generating initial codes, searching for

The transitional challenges and coping

themes among codes, reviewing themes, defining and naming themes, and producing the final report (9). The researchers analyzed with a research mentor who guided the researchers to ensure the empirical process of conducting the analysis. The coding and thematic analysis were done manually for better appreciation of the patterns and themes as the researchers fostered their

creativity and established a sense of ownership in the analysis. Data saturation was reached as themes emerged repetitively, and this led to the formulation of major themes and subthemes based from the prior codes.

The following table shows a sample of the thematic coding and analysis done in treating the data:

Table 1. Sample thematic coding and analysis of selected transcripts

Responses	Initial codes	Final theme
<i>"Most of us just started working here in just a few weeks or months. As a beginner in the psychiatric ward, I find a lot of flaws in terms of applying nursing care to my patients. I see the need to improve my skill set. I also need to grow professionally." (P3)</i>	<ul style="list-style-type: none"> • New to the work environment • Flawed nursing practice • Need to improve skills • Need to grow 	Nurses' recognition of the inability to patient care management
<i>"I thought that senior nurses should work closely with their juniors. Like me, I am new to the facility. I needed to seek help, especially when it is my first time managing a patient with such a condition. I talk to other nurses about what to do, but I feel ignored." (P2)</i>	<ul style="list-style-type: none"> • Senior nurses to work with new nurses • Need to seek help • Talk to another nurse • Felt ignored when asking help 	Need support from more experienced colleagues
<i>"I participate in research undertaking to hone my skills because whatever I do; it has to be based on the science and art of nursing practice. Research is important." (P3)</i>	<ul style="list-style-type: none"> • Participation in research • Practice-based in science and art of nursing • Importance of research 	Commitment to evidence-seeking practices
<i>"When patients trust you as his or her care provider, it will be very easy to handle him or her. I make sure that every time I engage with my patients, I establish rapport with them. It is the key to a successful therapeutic relationship. It is not easy at first, but it has to be established because it is really the key to caring for them." (P2)</i>	<ul style="list-style-type: none"> • Trust as a foundation of care • Establishing rapport • Therapeutic relationship 	Establishing therapeutic, professional nurse-patient relationship

Ethical considerations

The study was approved by the OLGC Research Evaluation Panel and thus was granted approval by the Evaluation Committee of the PTH headed by the director of the nursing service. Informed consent was obtained from the research participants using the World Health Organization (WHO) template. The participants were secured that all personal information gathered from them would remain anonymous. To protect their

identities, the researchers assigned pseudonyms to the hospital as "Psychiatric Training Hospital or PTH", and to the participants as "P1, P2, P3, P4, P5, and P6".

Trustworthiness

The study secured that this research was accurately measured and is reflective of the social reality of the participants (14). The triangulation process involved the presentation of the final report to the participants and sought their feedback

The transitional challenges and coping

responsibilities. Proper orientation to such regulations and policies in the institutions can prevent unwanted stresses at work.

"Being oriented to the policies is already a coping mechanism for me. Adherence to institutional policies is paramount." (P6)

Discussion

Generally, the findings showed that the challenges experienced by nurses during their transitory stage at work include emotional challenges, need for professional growth, nurses' recognition of the inability to patient care management, need support from more experienced colleagues, and experience/feel of shock in the encounter of mental patients. Meanwhile, the coping mechanisms that emerged after the thematic analysis revealed emotional responses, professional coping (i.e., reliance on the adequate support system, establishing the professional nurse-patient relationship, call to professional advancement, and commitment to evidence-based nursing practice), and institutional coping strategies.

Newly qualified nurses often feel overwhelmed by the challenges of the work environment and struggle to transition (15). Several stressors have been identified, and these stressors can be about consumer, family, and staff relationships, environment, and organization (16). The findings of the study support that newly qualified mental health nurses experience fear during their adaptation period, especially when they perceive mental healthcare users as dangerous and the working environment as risky, with little concern for their safety (15). The working environment of the nurses must be safe for them and their patients.

During the transition, nurses require opportunities to develop both competence and confidence in their ability to practice independently (15). New nurses in a mental health environment learn from accepting responsibility, taking on challenges, and making adjustments to develop confidence as mental healthcare professionals (15). Efforts to cope could be cognitive and behavioral (17). All stakeholders should take

proactive measures to alleviate challenges among nursing staff (18) as work correlates to stress (19).

Emotional challenges in newly hired psychiatric nurses are attributed to the challenging and potentially high-stress setting (16). Considering the behavioral patterns exhibited by patients with schizophrenia, newly hired nurses need to be more prepared in the care of their patients. Psychiatric nursing requires nurses to adapt despite the wide range of stressful events, evolving from the care of violent, aggressive patients, recurrent relapse to poor prognosis of mental disorders (20). In a study, psychiatric nurses also had moderate levels of work-related stress and depression and exhibiting different coping mechanisms (20).

The limited exposure during their baccalaureate degree was cited to be contributory to their current flawed practice. This implies that the knowledge and skill required of mental health nurses to provide physical health care need to be increased (21). From their experiences, some have forgotten that the patients also need adequate physical care; hence, the improvements of the nursing education (22) curriculum are vital to prepare future nurses.

The current study also highlights that institutions should employ expert nurses who can guide mental health nurses to provide physical health care (21) in this regard. In a study, it has been proposed that institutions should employ expert nurses who can guide mental health nurses to provide physical health care (21).

The participants of the current study revealed their therapeutic practices to establish therapeutic relationships. These relationships are powerfully influenced by individual and organizational factors (23). Necessarily, caring nurse-patient relationships in mental health settings are critical components in helping patients recover (24).

Senior nurses, the more experienced ones, need to extend assistance to newly hired nurses to meet desired therapeutic ends. Imparting the skills and experiences of

senior nurses to novice nurses and their presence alongside other nurses on work-shifts can be useful in promoting vigilance (25).

Despite the challenges, the participants are committed to their jobs. The motivation to pursue a career in psychiatric nursing, especially for the new psychiatric nurses, is rooted in the provision of high quality, meaningful placements (26). Providing care to patients with psychiatric disorders and working in a challenging environment lead to many problems among psychiatric nurses (27). Addressing stigma and more exceptional education and support for nurses to pursue a mental health career (26) must also be enforced. It is the moral and professional responsibility of the institution to protect them (27).

In terms of coping mechanisms, these are essential (28) to psychiatric nurses. These enable newly hired nurses to counteract their transitional challenges. Implementing programs aimed at teaching nurses how to deal with stress at work and improving their coping mechanisms and problem-solving skills are recommended (20). Administrative and collegial supports are also vital among newly hired psychiatric nurses. These are critical because support from supervisors and working environments can reduce job stress and negative emotions toward patients (29). To keep newly hired psychiatric nurses committed and motivated, there is a need to satisfy their various motivations to stay despite the challenges. A recent study suggested that the government and hospital administrators should consider ways to improve income, reduce work hours, and promote the psychiatric nursing specialty in ways that increase the public's respect for it, increase awareness of medical liability insurance coverage, and protect nurses from patients' violence (30).

Moreover, in the current study, the results imply the resilience of the newly hired psychiatric nurses as a manifestation of their coping mechanisms. Newly hired psychiatric nurses revealed the temporality of their challenges, which speaks much of their resilience as they care for their patients.

Resilience has positive correlations with hardiness, self-esteem, life and job satisfaction, and negative correlations with depression and burnout (FOSTER). Through a resilience program, improvement in coping self-efficacy and capacity to regulate thoughts and emotions (FOSTER) is most likely to develop.

Implications of the Study

The findings of this study reverberate the importance of preparing nurses to work in mental health facilities. Like any other field of nursing practice, psychiatric and mental health nursing, as a highly specialized field of nursing, requires the right preparations in terms of knowledge, skills, and attitudes. Adequate and reasonable training and exposure to mental health facilities where they can observe and practice during their baccalaureate degree must be warranted to help them adjust to the actual world of work should they desire to practice mental health nursing. Good training and exposure during their RLE can best prepare them to minimize their concerns relative to their transitory stage. Secondly, the findings also emphasize the need to equip nurses continuously to become more competent and compassionate in psychiatric and mental health nursing. The need for newly hired nurses for continuous growth and development is crucial to overcome their emerging challenges, especially during the transitory stage of their careers. Lastly, the results of the study highlight the reliance on adequate support systems in the workplace for newly hired nurses to overcome challenging situations. A mentoring program may be institutionalized and facilitated were more experienced nurses' guide and assist newly hired nurses in delivering quality patient care. Likewise, a sustainable resilience program may also be established to help newly hired nurses confront their challenges with endurance.

Conclusion

Caring for patients with mental disorders has never been easy for newly hired nurses. The concern can pose a significant effect on

12. Benner P. From novice to expert. Menlo Park. 1984.
13. Patton, MQ. Qualitative Research & Evaluation Methods: Integrating Theory and Practice (4th ed.). USA: Sage Publications, Inc.2015
14. Maher C, Hadfield M, Hutchings M, de Eyto A. Ensuring rigor in qualitative data analysis: A design research approach to coding combining NVivo with traditional material methods. *International Journal of Qualitative Methods*. 2018 Jul 9;17(1):1609406918786362.
15. Mabala J, van der Wath A, Moagi M. Newly qualified nurses' perceptions of working at mental health facilities: A qualitative study. *Journal of psychiatric and mental health nursing*. 2019 Jun;26(5-6):175-84. doi:10.1111/jpm.12525.
16. Foster K, Roche M, Delgado C, Cuzzillo C, Giandinoto JA, Furness T. Resilience and mental health nursing: An integrative review of international literature. *International journal of mental health nursing*. 2019 Feb;28(1):71-85.
17. Song Y, Ebenezar M, Haastrup K. Coping mechanisms of mental health nurses in work related stress situations: Literature Review. 2018. Availale from: https://www.theseus.fi/bitstream/handle/10024/161218/Moteka_Ebenezar_Haastrup_Kolade_YuZhen_Song_Pdf.pdf?sequence=2&isAllowed=y
18. Haslinda A, Tyng CL. Job stress and coping mechanisms among nursing staff in a Malaysian private hospital. *International Journal of Academic Research in Business and Social Sciences*. 2016 May;6(5):471-87.
19. Sari DH, Erwina I. Relationship between Job Stress and Coping Mechanism in Psychiatric Nurse in National Mental Health Hospital. *Elevate: The International Journal of Nursing Education, Practice and Research*. 2018 Jul 1;1(1):76-9.
20. Hasan AA, Elsayed S, Tumah H. Occupational stress, coping strategies, and psychological-related outcomes of nurses working in psychiatric hospitals. *Perspectives in psychiatric care*. 2018 Oct;54(4):514-22. doi:10.1111/ppc.12262
21. Çelik Ince S, Partlak Günüşen N, Serçe Ö. The opinions of Turkish mental health nurses on physical health care for individuals with mental illness: a qualitative study. *Journal of Psychiatric and Mental Health Nursing*. 2018 May;25(4):245-57. doi:10.1111/jpm.12456
22. Harrison CA, Hauck Y, Ashby R. Breaking down the stigma of mental health nursing: A qualitative study reflecting opinions from western australian nurses. *Journal of Psychiatric and Mental Health Nursing*. 2017 Sep;24(7):513-22. doi:10.1111/jpm.12392.
23. Pazargadi M, Fereidooni Moghadam M, Fallahi Khoshknab M, Alijani Renani H, Molazem Z. The therapeutic relationship in the shadow: Nurses' experiences of barriers to the nurse–patient relationship in the psychiatric ward. *Issues in Mental Health Nursing*. 2015 Jul 3;36(7):551-7. doi:10.3109/01612840.2015.1014585
24. Valente SM. Managing professional and nurse–patient relationship boundaries in mental health. *Journal of psychosocial nursing and mental health services*. 2017 Jan 31;55(1):45-51. doi:10.3928/02793695-20170119-09
25. Vahidi M, Ebrahimi H, Areshtanab HN, Jafarabadi MA, West S, Foong A, Cleary M. Providing a safe environment in a psychiatric ward: Nurses and vigilance in Iran. *Issues in mental health nursing*. 2019 Jan 2;40(1):65-72. doi:10.1080/01612840.2018.1463326
26. Penman J, Martinez L, Papoulis D, Cronin K. Voices from the Field: Regional Nurses Speak About Motivations, Careers and How to Entice Others to Pursue Mental Health Nursing. *Int J Nurs Educ Scholarsh*. 2018;15(1):/j/ijnes.2018.15.issue-1/ijnes-2017-0056/ijnes-2017-0056.xml. Published 2018 30 January. doi:10.1515/ijnes-2017-0056
27. Zarea K, Fereidooni-Moghadam M, Baraz S, Tahery N. Challenges encountered by nurses working in acute psychiatric wards: A qualitative study in Iran. *Issues in Mental Health Nursing*. 2018 Mar4;39(3):244-50. doi:10.1080/01612840.2017.1377327
28. Tsaras K, Daglas A, Mitsi D, Papatathanasiou IV, Tzavella F, Zyga S, Fradelos EC. A cross-sectional study for the impact of coping strategies on mental health disorders among psychiatric nurses. *Health psychology research*. 2018 May 8;6(1). doi:10.4081/hpr.2018.7466
29. Matsumoto Y, Yoshioka SI. Factors Influencing Psychiatric Nurses' Job Satisfaction Levels: Focusing on Their Frequency of Experiencing Negative Emotions Toward Patients and Support at Their Workplaces. *Yonago Acta Medica*. 2019;62(4):293-304. doi:10.33160/yam.2019.11.006
30. Zhou H, Jiang F, Rakofsky J, Hu L, Liu T, Wu S, Liu H, Liu Y, Tang YL. Job satisfaction and associated factors among psychiatric nurses in tertiary psychiatric hospitals: Results from a nationwide cross-sectional study. *Journal of Advanced Nursing*. 2019 Dec;75(12):3619-30. doi:10.1111/jan.1420.